

# Investigation of atoms in electric fields: the Stark effect

Masud Karim Sarkar

Department of Physics , Bidhan Chandra College, Rishra, Hooghly, West Bengal, India (Affiliated to Calcutta University)

## ARTICLE DETAILS

### Article History

Published Online: 07 September 2018

### Keywords

Stark effect, electric, atoms

## ABSTRACT

A traditional model of the hydrogen particle in a static electric field is considered, basing upon the work [Hooker A. et al, Phys. Fire up. A, 55 (1997) 4609]. In that work the electrons should move along Kepler circles around the core, while communicating with the outer field. That traditional model imitates very well the genuine energy move of the direct Stark impact. The concurrence with the subsequent request impact is poor. It is appeared here that the outcomes for the quadratic Stark impact might be impressively improved in the event that the electrons are as yet permitted to move along traditional directions, yet whose starting conditions are factually examined from a circulation taking after the quantum mechanical one.

## 1. Introduction

The spectrum of the hydrogen atom in frail opposite and equal electric and attractive fields is processed and broke down with stipend for the diamagnetic connection. In either case the energy spectrum of the profoundly energized states as a component of the electric field power for a fixed attractive field separates into three subjectively various districts. A characteristic clarification of this impact is found in a quasiclassical examination of three fold recursion relations to whose arrangement both issues can be decreased. The reason for this parting of the spectrum lies in the rebuilding of the states, when the connection between the electric-and attractive field powers is changed, therefore of the appearance or vanishing of successful likely boundaries. Those parts of the quasiclassical estimate for the three fold recursion relations which are essential for the examination of the two issues and which have up to this point not been talked about in the writing are thought of.

As of late there has been an upsurge in interest in the issue of the portrayal of the Rydberg states in a frail attractive field. This is expected above all else to the revelation of the dramatic decline of the energy-level parting at the quasicrossing point as the central quantum number  $n$  increments. This conduct of the parting was found first in a mathematical calculation<sup>1</sup> and afterward tentatively.<sup>1</sup> To clarify this wonder, Zimmerman et al.<sup>1</sup> and Delande and Gay<sup>2</sup> construed the presence of an inexact concealed balance. A reliable depiction of the surmised balance and a subjective clarification of the dramatic decline of the parting are introduced in Ref. 3, where it is demonstrated that for the hydrogen atom in a powerless attractive field  $H$  the factors of the Schrodinger condition with terms up to the fourth request in  $H$  are detachable in elliptic round and hollow arranges on a circle in four-dimensional energy space. In Ref. 4 an explanatory articulation is gotten for the size of the parting at the quasicrossing point. Different qualities of the spectrum are likewise considered in Ref. 4; specifically, it is found there that the lower energy levels in a multiplet with fixed estimations of  $n$  and  $(m$  is the azimuthal quantum number) are roughly twofold savage for the situation when  $n > m$ . The parting of these doublets is registered in Ref. 5. The partition of the factors in force space is talked about in Refs. 4 and 6. In Refs. 7 and 8 a

few variations of the quasiclassical estimate are viewed as which comparable to the one are utilized in Ref. 3. In Refs. 4-7 surmised investigative articulations are gotten for the lower and upper energy levels in a given multiplet

The utilization of old style models to examine quantum mechanical frameworks is to a great extent utilized in minute material science. For huge quantum numbers it is advocated on the premise of the correspondence rule. Additionally at little quantum numbers its utility isn't immaterial since it permits in any event to get a subjective knowledge of the elements of the framework; in the best cases, it is even conceivable to extricate quantitative outcomes in a simple manner and with a decent exactness. A few models are the dissipating of electrons from a particle, and the treatment of vivacious particle molecule impacts. Old style model are clearly preferred over their quantum-mechanical partners from the mathematical purpose of see. In those situations where additionally the quantum issue is feasible, the traditional picture is significant since it permits to obviously envision the current issue. Consequently the correspondence between the traditional and quantum mechanical depictions is as of now a very much considered subject. Recently it has been proposed an absolutely traditional view to the Stark impact in hydrogen: the field should collaborate with traditional electrons pursuing Kepler circles around the core. Following this work, in ref. the energy shifts for the straight and quadratic Stark impact have been registered. The arrangement between the traditional outcome and the specific quantum mechanical one is magnificent for the direct impact yet isn't useful for the quadratic impact, which just asymptotically for enormous quantum numbers approaches the right worth. The motivation behind the current paper is to show that, inside a simply traditional formalism fundamentally the same as that of, the quadratic Stark impact might be replicated precisely: it is sufficient to loosen up the limits on the electron direction and receive a factual methodology: here, the electron is as yet demonstrated as a traditional molecule following Kepler circles however the underlying conditions are picked up from a distribution subject to certain rules. It will be shown that a much better agreement is obtained by this slightly more sophisticated approach. Atomic units will be used throughout this work unless explicitly stated.

**2. The classical model**

Also known, the investigation of the Stark impact is effortlessly acted in explanatory organizes, where the hydrogen molecule is characterized by the quantum numbers  $n, n_1, n_2, m$ . We are thinking about little fields, for which a perturbative approach is sufficient they actually are acceptable quantum numbers. I will first quickly sum up the treatment of the primary request Stark impact, which will be helpful additionally to present the fundamental ideas utilized later. First of all, let us review a few outcomes expressed in. In that work a correspondence has been done between the quantum mechanical administrators and their classical partners:

- to each set of qualities  $n_1, n_2, n, m$  there compares a bunch of classical Keplerian circles of the electron which are commonly circular, with flightiness  $\epsilon = p_1 - l_2/n_2$  ( $l$  is rakish quantum number in round directions, and the classical precise force). At the point when time–found the middle value of over a time of revolution, the oval yields a non zero mean electric dipole second:

$$\langle d \rangle = 1T \int_0^T z dt = 3/2 n^2 \epsilon .$$

- The Lenz vector  $A = p \times L - r/|r|$  is classically a conserved quantity, as is its  $z$ -component in quantum mechanics:

$$A_z = (n_1 - n_2)n .$$

$A$  is related to the electron orbit by  $|A| = \epsilon$ . Since, besides this, in it was shown that  $d$  and  $A$  point towards the same direction, one may identify them through

$$d = (3/2)n^2 A .$$

The energy shift in presence of an electric field  $F$  is calculated from

$$\Delta E^{(1)} = -d \cdot F .$$

It is immediate to see that this result agrees with the energy shift obtained using quantum mechanics provided.

$$\Delta E^{(1)} = -32n(n_1 - n_2)|F| ,$$

In it was checked by mathematically incorporating the Kepler circles in presence of an electric field and with introductory conditions viable with the quantized worth that the classical energy shifts from its unperturbed estimation of the amount .

The quadratic impact is of significance when  $n_1 = n_2$ , in which case the direct term disappears. This implies in the language of –that the mean dipole second disappears, and this happens when the circles are roundabout and lying on a plane opposite to the field pivot. The electric field may prompt a dipole second by moving the electron and the core as for each other. The energy move for the quadratic Stark impact is characterized by

$$\Delta E^{(2)} \equiv -12\alpha F^2 ,$$

$\alpha$  being the polarizability.  $\alpha$  is related to the induced dipole moment by

$$\alpha F = d .$$

The quantum mechanical value of  $\alpha$  for states with  $n_1 = n_2$  and  $m = n - 1$  is

$$\alpha_{QM} = n^4 4(4n^2 + 9n + 5) .$$

In a straightforward estimation is utilized to figure the classical estimation of  $\alpha$ , acl: be  $r$  the span of the round circle of the electron, and  $\delta z$  the move along the course of  $F$  between the electron and the core instigated by the outside field. For little estimations of  $\delta z$  we may inexact the Coulomb power on the electron as  $\delta z/r^3$ , which adjusts the power applied on the electron by  $F$  when

$$F = \delta z r^3 .$$

The induced average dipole moment is  $d = \delta z$  and, from eq. (9) and the definition of  $\alpha$

$$\alpha_d = r^3 = n^6$$

Since for round Bohr circles  $r = n^2$ .

Regardless of whether this is an extremely disentangled model, it was checked in that it depicts precisely the classical framework: the Hamilton's conditions for the electron were mathematically unraveled in presence of the electric field. The electron energy was resolved from its position and force. The mathematical outcomes were found to concur for  $n \rightarrow \infty$ ; the more noteworthy disparities are for little  $n$ s: for instance, when  $n = 1$  the proportion is acl/ $\alpha_{QM} = 2/9 \approx 0.22$ .

**3. Conclusions**

In ref. it is commented that the primary inspiration of their work was to show that some quantal outcomes might be recuperated, at any rate somewhat, inside a simply classical structure, given just that the underlying conditions be picked as viable as conceivable with the laws of quantum mechanics. This paper follows precisely that line of thought, being a refinement of that work in that the underlying conditions have been picked in a more right way, however as yet staying inside a classical portrayal of the framework. One may ponder on the off chance that the value paid to have this more noteworthy exactness is excessively high, since we utilized careful wave–capacities, which is identical to understand the quantum mechanical issue. It isn't thus, since huge upgrades are acquired utilizing no matter which appropriation, even that of eqn. which depends on simply classical contemplations. The other two conveyances have been picked just to give the peruser a correlation. This is significant for every one of these cases where extricating the wave work is too troublesome and one is compelled to resort to approximations. As effectively called attention to in, it very well may merit investigating in some of such circumstances.

**References**

- [1] Borodin V. M., Kazansky A. K. and Ochkur V. I., *J. Phys. B: At. Mol. Opt. Phys.*, 25 (1992) 445.
- [2] Rost J.-M. , *Phys. Rev. Lett.*, 72 (1994) 1998; Rost J.-M. , *J. Phys. B:At. Mol. Opt. Phys.*, 27 (1994) 5923.
- [3] Ivanovski G., Janev R. K. and Solov'ev E. A., *J. Phys. B: At. Mol. Opt. Phys.*, 28 (1995) 4799.
- [4] Abrines R.A. and Percival I. C., *Proc. Phys. Soc.*, 88 (1966), 861; *ibid*873; Bransden B. H. and McDowell M. R. C., *Charge Exchange and the Theory of Ion-Atom Collisions* (Clarendon Press, Oxford), 1992, ch. 8.
- [5] Ostrovsky V. N., *J. Phys. B: At. Mol. Opt. Phys.*, 28 (1995) 3901.
- [6] Hezel T. P., Burkhardt C. E., Ciocca M. and Leventhal J. J., *Am. J. Phys.*, 60 (1992) 324; Hezel T. P., Burkhardt C. E., Ciocca M., He L.-W., and Leventhal J. J., *Am. J. Phys.*, 60 (1992) 329.
- [7] Hooker A., Greene C. H., and Clark W., *Phys. Rev. A*, 55 (1997) 4609.
- [8] Galindo A. and Pascual P., *Quantum Mechanics* (Springer-Verlag), 1991, ch. 10.
- [9] Cohen J. S., *J. Phys. B: At. Mol. Phys.* , 18 (1985) 1759.