

Effect of parallel resistance and plyometric training on coordinative abilities among university level cricket players

¹S.Selvakumar & ²Dr.P.Sivaraman

¹Ph.D Research Scholar, Department of Physical Education, Annamalai University, Annamalai Nagar-608002, Tamilnadu (India)

²Assistant Professor, Department of Physical Education, Annamalai University, Annamalai Nagar-608002, Tamilnadu (India)

ARTICLE DETAILS

Article History

Published Online: 10 December 2018

Keywords

Parallel Resistance training, Coordinative abilities, Orientation ability, Cricket players

ABSTRACT

For the purpose of the study sixty (60) were randomly selected as subjects and the age was ranged between 18 and 25 years. The selected subjects were randomly assigned into two equal groups such as training group (TG) and the control group (CG) for the strengths of fifteen (n = 30) each. Experimental group underwent respective Parallel Resistance training programme on coordinative abilities for twelve (12) weeks for four to five days. The control group did not involve in any special training apart from their regular activities. The coordinative abilities were taken as a criterion variable for the present study and it was measured by orientation ability test. Analysis of covariance (ANCOVA) was used to analyse the collected data. The results revealed that the Parallel Resistance training made significant improvement on the coordinative abilities among of the selected subjects. The level of confidence was fixed at 0.05 in all cases.

1. Introduction

The game of cricket has historically been known as "the gentleman's game". Until about three decades ago cricketers were certainly not the fittest athletes on the planet. Often it was remarked that cricket is physically an easy game which requires one to stand on the field for most of the day and requires little running, jumping or strength (Kushwah, 2012). The highly developed levels of fielding in the modern times require a player to have strong shoulders and arms to make direct hits at the stumps. The modern player is leaner, stronger and far more athletic. For cricket enthusiasts there is nothing to match the meaningful contests and excitement generated by the game's subtle shifts in play (Hardayalsingh, 1984).

The British brought cricket to India in the early 1700 with the first cricket match played in 1721. In 1848, the Parsi community in Bombay formed the Oriental Cricket Club, the first cricket club to be established by India Ltdns. After slow beginnings, the Europeans eventually invited the Parsis to play a match in 1877. By 1912, the Parsis, Sikhs, Hindus and Muslims of Bombay played a quadrangular tournament with the Europeans every year. In the early 1900s, some Indians went on to play for the England cricket team. Some of these, such as Ranjitsinhji and KS Duleepsinhji were greatly appreciated by the British and their names went on to be used for the Ranji Trophy and Duleep Trophy – two major first-class tournaments in India. In 1911 an Indian team went on their first official tour of the British Isles, but only played English county teams and not the England cricket team.

Resistance training is a form of strength training in which each effort is performed against a specific opposing force generated by resistance i.e. resistance to being pushed, squeezed, stretched or bent (Syed et al, 2006). Exercises are isotonic if a body part is moving against the force. Exercises are isometric if a body part is holding still against the force. Resistance exercise is used to develop the strength and size of skeletal muscles. Property performed, resistance training can

provide significant functional benefits and improvement in overall health and well being .

Many years coaches and athletes have sought to improve power in order to enhance performance. Throughout this century and no doubt during the times before, jumping, bounding and hopping exercises were especially used in various ways to enhance athletic performance. In recent years this distinct method of training for power or explosiveness has been termed Plyometrics (Kiranpal Singh, 2004). Plyometrics by definition is, a type of exercise using explosive movements to develop muscular power, bounding, hopping and jumping.

Coordinative ability means an ability to quickly and purposefully perform difficult spatio-temporal movement structures. Within this context, coordination abilities are understood as an externally visible manifestation of the control and regulation processes of the motor activity of the central nervous system (Simonek and Jaroslav, 2009). The complex of coordination abilities consists of a group of basic coordination abilities.

2. Methodology

To achieve the purpose sixty(60) cricket players were selected as subjects from Annamalai University, Annamalai nagar Tamil Nadu India, the age of the selected subjects was 18-25 years. The subjects were divided in to two groups namely physical training (n=30) and control (n=30). The physical training group underwent training for 60 min/4 days/12 weeks including warming up and cooling down exercises.

Experimental training group underwent Parallel Resistance training programmes for 12 weeks for four to five days per week. The control group did not involve in any special training apart from their regular activities. The coordinative abilities were taken as a criterion variable for the present study. The collected data were statistically examined by analysis of

covariance (ANCOVA). The confidence level was fixed at 0.05 levels, which is appropriate to the present study.

3. Result discussion and Findings

TABLE – I.1
ANALYSIS OF COVARIANCE OF PRETEST POST TEST AND ADJUSTED POST TEST
ON ORIENTATION ABILITY OF DIFFERENT GROUPS

Test	EX.1	EX.2	EX.3	CG	SV	SS	DF	MS	F
PRETEST	11.41	11.56	11.78	11.30	B	1.92	3.00	0.64	2.47
					W	14.56	56.00	0.26	
POST TEST	10.28	10.11	9.63	11.75	B	37.69	3.00	12.56	47.98*
					W	14.66	56.00	0.26	
ADJUSTED	10.28	10.10	9.62	11.75	B	34.56	3.00	11.52	43.23*
					W	14.66	55.00	0.27	
MEAN DIFF.	-1.13	-1.46	-2.15	0.45					

* Significant at .05 level of confidence.

SV-Source variances, SS-Sum of squared, df-degrees of freedom, MS-Means square.


(The table values required for significance at .05 level of confidence for 3 and 56 and 3 and 55 are 2.77 respectively).

TABLE – I.2
Scheffe’s post hoc test mean differences on Orientation ability among four groups

EXP.1	EXP.GP 2	EXP.GP.3	CONTROL	MD	CI
10.28	10.10			-0.18	0.54
10.28		9.62		-0.66*	0.54
10.28			11.75	1.47*	0.54
	10.10	9.62		-0.48	0.54
-	10.10		11.75	1.65*	0.54
		9.62	11.75	-2.13*	0.54

* Significant at 0.05 level of confidence.

Fig. (i)



4. Discussion

The result of the table I.1 shows the analyzed data on Orientation ability. The pre-test means of Orientation ability were 11.41 for resistance training group, 11.56 for plyometric training group, 11.78 for combined group and 11.30 for control group. The obtained “F” ratio of 2.47 was lesser than the table F-ratio 2.77. Hence the pre-test was not significant at 0.05 level of confidence for the degrees of freedom 3 and 56.

The post-test means of Orientation ability were 10.28 for resistance training group, 10.11 for plyometric training group, 9.63 for combined group and 11.75 for control group. The obtained “F” ratio of 47.98 was greater than the table F-ratio 2.77. Hence the post-test was significant at 0.05 level of confidence for the degrees of freedom 3 and 56.

The adjust post-test means of Orientation ability were 10.28 for resistance training group, 10.10 for plyometric training group, 9.62 for combined group and 11.75 for control

group. The obtained "F" ratio of 43.23 was greater than the table F-ratio 2.77. Hence the adjust post-test was significant at 0.05 level of confidence for the degrees of freedom 3 and 56.

Since, four groups were compared, whenever the obtained 'F' ratio for adjusted post test was found to be significant, the Scheffe's test to find out the paired mean differences and it was presented in Table I.1.

Table - I .2 shows the Scheffe's post-hoc test results. The ordered adjusted final mean difference for orientation ability of isolated and combined resistance and plyometric training groups and control group were tested for significance at 0.05 level of confidence against confidential interval value.

All the training groups namely resistance training, plyometric training and combined training and control group were 1.47, 1.65 and 2.13 respectively and it was seen to be greater than the confidential interval value of 0.54. Hence all the training groups were better than the control group.

The mean differences between resistance training and combined training was 0.66 it was greater than confidence interval 0.54, hence there was a significant difference between these groups. Resistance training group and plyometric training group was 0.18 and plyometric training and combined training group was 0.48, it was found to be lesser than confidential interval value of 0.54, hence these groups were equal in improving the orientation ability. The mean values of orientation ability are shown graphically in **Figure. (i)**Above.

5. Findings

Sharma and Gangwar (2014) conducted a study was to investigate difference of coordinative abilities between footballer and hockey male players. To attain this study, 30 (15 footballer and 15 hockey players) represented inter-university tournament with age ranging from 17 to 25 years were randomly selected from Lakshmbai National Institute of Physical Education, Gwalior. The statistical technique employed for this study was independent 't'-test at 0.05 level of significance. As per the statistical analysis significant difference was found between football and hockey players on

the sub-variables i.e. reaction ability, orientation ability and differentiation ability, However insignificant difference was found on sub-variable i.e. rhythmic ability at 0.05 level of significance.

Bal and Sandhu (2014) conducted a study is to find out the significant differences of psychomotor abilities, among cricket players of different level of achievement. For the purpose of present study, One hundred seventy seven (N=177), Male District, State and National Level Cricket Players between the age group of 21-25 years (Mean $\frac{1}{2}$ SD: age 22.89 $\frac{1}{2}$ 1.76 years, height 176.04 $\frac{1}{2}$ 4.18 cm, body mass 73.76 $\frac{1}{2}$ 4.63 kg) were selected. The subjects were purposively assigned into three groups: Group-A: District level cricket players (n=80), Group-B: State Level Cricket Players (n=65) and Group-C: National level cricket players (n=32). The statistical package for the Social Sciences (SPSS) version 14.0 was used for all analyses. In all the analyses, the 5% critical level (p 0.05) was considered to indicate statistical significance. The differences in the mean of each group for selected variable were tested for the significance of difference by One-way Analysis of Variance (ANOVA). For further analysis Post-Hoc Test (Scheffe's Test) was applied. The results revealed significant differences between found among district, state and national level cricket players on the sub-variables; muscular strength, muscular power, muscular endurance, running speed, running agility, jumping ability, throwing ability, flexibility and balance.

6. Conclusion

The result of the study revealed that the Parallel Resistance training and Plyometric training has significant improvement on Coordinative abilities among University level cricket players. It was also concluded that the parallel resistance training method and plyometric trainings are one of the best training methods for improving orientation abilities as well as the physical fitness components among university level cricket players.

References

1. Deepak Singh Kushwah, Comparison of Differentiation Ability and Rhythm Ability of Basketball and Handball Players, International Journal of Health, Physical Education and Computer Science in Sports, Vol 7, Issue 1,(2012) p. p. 175-177.
2. Hardayal Singh, Sports Training - General Theory and Technique, (Patiala: 1991),.p.177.
3. I.Syed, A.Kanniyan, H.A. Moslim and M.A. Syed, Coordinative Ability Variables among Tennis and Squash Men Players – A Comparative Study, British Journal Sports Medicine, (2014) p. p. 48:663.
4. Jaromir Simonek and Jaroslav Brodani, Structure of Coordination Abilities and Prediction of Coordination Performance in the Selected Sports, Sport Science Review, Vol. 28, Issue 5- 6 (2009).
5. Kiranpal Singh, Comparison of Selected Coordinative abilities among Sportsmen belonging to contact semi-contact and non-contact sports, Unpublished Thesis Doctor of Philosophy, Lakshmbai National University of Physical Education, Gwalior (M.P.),2004.
6. Sandeep Sharma and Narendra Gangwar, Comparison of selected Coordinative abilities between football and hockey male players, IJMESS, P-ISSN: 2278-0793 and E-ISSN: 2321- 3779, Vol. 3 No. 1 (March, 2014).
7. Baljinder Singh Bal and Ranjeet Singh Sandhu, An Exploratory Study of Psychomotor Abilities among Cricket Players of Different Level of Achievement, Res. of Physical Education Sci., ISSN-2320-9010, Volume 2, Issue (8), August (2014) p.p.- 1-19