

Study the Framework, Determination and Assemblage of the Fish from the Ganga River

Neelam

M.Sc. in Geography H.N. 1432, SEC -2, Bhadurgarh, Jhajjar Haryana (India)

ARTICLE DETAILS

Article History

Published Online: 13 March 2019

Keywords

Wellspring of Sustenance, Change of Hydrology and Presentation of Intriguing Species, Cypriniformes and Cyprinid

*Corresponding Author

Email: neelamgeo2018@gmail.com

ABSTRACT

Fisheries are a noteworthy wellspring of sustenance, recreation and salary for mankind and fishers network universally. The stream Ganges is the biggest waterway in India and the fifth longest on the planet. Albeit, numerous studies on fish ecology and systematic have been directed generally to improve fisheries however fish decent variety and their dispersion design from preservation perspective have never been enough tended to in the Ganges. The Ganga waterway fauna and vegetation is undermined by anthropogenic exercises and coming about water contamination, amassing of substantial metals, eutrophication, damming, and change of hydrology and presentation of intriguing species. Fish gatherings are an essential segment of amphibian ecosystems. Present examination was attempted to consider the Determination, structure and wealth of fishes from the Ganga waterway (site 1: Kanpur, site 2: Allahabad and site, 3: Varanasi segment), India. The Ganga River is a spine of Indian fishery. The examples were gathered month to month amid the period June 2011 to May 2013. Sanctioned correspondence analysis (CCA) showed that hub 1 and 2 represented 67% and 33% fluctuation for species and environmental connection, individually. Structures of the fish gathering of the Ganga at Kanpur to Varanasi harbors of 102 fish species have a place with 8 requests and 28 families. 74, 89 and 82 fish species were recorded at Kanpur, Allahabad and Varanasi destinations, individually. Cypriniformes and Cyprinid were the most rich fish species request and family in all destinations.

1. Introduction

Freshwater fishes are critical and esteemed property for money, human sustenance, game and adornment. Overexploitation happens far and wide with the utilization of increasingly more refined fishing hardware, and the abatement of many fish stocks has been archived because of growing fisheries. Illicit fishing utilizing explosive, pesticides, electro-fishing, and so on is likewise real dangers to angle biodiversity everywhere throughout the world.

Fish gatherings are recognized as responsive pointers of living space corruption, natural condition debasement, environmental defilement, and in general ecosystem profitability. Freshwater fishes are the most jeopardized vertebrate group with an anticipated annihilation rate of multiple times that of earthly fauna and multiple times that of marine warm blooded animals. The world natural fishery systems are crumbling as an immediate aftereffect of overfishing and overcapacity of fishing armadas. Biodiversity is basic for adjustment of ecosystem, insurance of by and large environmental quality for understanding natural worth of all species on the earth. Biodiversity of fishes are enduring step by step in 21st century. Primary districts are accessibility of water in waterway/stream (for haven), water deliberation, enterprises and private use, territory obliteration and defragmentation, contamination level, presentation of outsider/extraordinary species and effects of worldwide atmosphere changes uniquely precipitation. Circulation examples of life forms are constrained by dispersal instrument, historical components (interfacing pathways, dispersal boundaries) and resistance to environmental elements.

Biodiversity is the amount, assortment and appropriation crosswise over organic scales running through hereditary qualities and life types of populaces, species, networks and ecosystems. Biodiversity influences the limit of living systems to react to changes in the environment, supports ecosystem work and gives the ecosystem goods and services that help human prosperity (e.g., supplement cycling, clean water).. This area (Kanpur to Varanasi segment) of the stream is center stretch which is most imperative for fisheries and human obstruction. In any case, no data is accessible on Canonical Correspondence Analysis (CCA) in the Ganga River particularly from Kanpur to Varanasi area, India.

At all out stretch, Cypriniformes request was shared 49 species, trailed by Siluriformes 26 species and Perciformes 17 species. Requests Clupeiformes shared 5 species. Plenitude was ruled by *Eutropiichthysvacha* contrasted with *Clupisomagarua* and *Sperataseenghala*. As indicated by wealth, *Cyprinus carpio* var. *communis* (9.64%) and *Oreochromis niloticus* (9.19%) were intensely trespasser in the Ganga River. Outlandish species is disturbing for indigenous species biodiversity. *C. carpio* var. *communis* and *O. niloticus* are every now and again recorded in the Ganga River. All out hardness, alkalinity and broke up oxygen were in charge of the nearness of catla, rita and *Sperataaor*, while *Labeocalbasu*, *Cyprinus carpio* and *Cirrhinus mrigala* favored nitrate, phosphate and all out disintegrated strong for their bounty. *Oreochromis niloticus* favored high organic oxygen request and lead while Zn and Sulfate were in charge of wealth of *L. rohita*. For protection perspective *C. carpio* var. *communis* and *O. niloticus* species ought to be observed in the Ganga River. The two species are extremely destructive for fish biodiversity in

the Ganga River. Fish gathering and their plenitude know the soundness of ecosystem.

2. Objective of the Study

The target of the present examination was to give Canonical Correspondence Analysis (CCA) of the Ganga River at Kanpur to Varanasi area, expecting to contribute a superior learning to structure of the fish collection and plenitude of economically vital fishes from the Ganga River and an apparatus for protection arranging of amphibian environments in this district.

3. Material and methods

The examples were gathered month to month amid the period August 2012 to July 2013 from the three locales of the Ganga stream in particular Kanpur (Latitude-26° 27' 16" N, Longitude-80° 20' 58"), Allahabad (Latitude-25° 45' 27" N, Longitude-81° 59' 31") and Varanasi (Latitude-25° 19' 01" N, Longitude-82° 58' 15"). Present stretch is around 370 km.

Human exercises and mechanical influent greatest revealed in these destinations of the stream. The Ganga River is a blessed waterway of India and has been proclaimed as a national stream by the administration of India. The Ganga is an enduring waterway which begins as a stream called "Bhagirathi" from Gaumukh (Himalaya) in the Gangotri ice sheet at 30° 55' N, 79° 7' E, and somewhere in the range of 4100 m above mean ocean level. Ganga waterway bowl is the biggest stream bowls in India and the fourth biggest on the planet, with a bowl (catchment region) covering 8, 61,404 sq km. It has an all out length of 2525 km with two nations (India and Bangladesh). It is spine for water system, agriculture, and industrials reason and fisheries perspective.

Sanctioned correspondence analysis and related strategy has discovered wide-spread use in amphibian sciences. Standard Correspondence Analysis (CCA rendition 4.5) was utilized to inspect the components reaction for the bounty of fish wealth.




Figure 1 showing the sampling sites.

The gathered examples were protected in 10% formalin and conveyed to the research center for further investigation. The fish was recognized utilizing Day, Talwar and Jayaram books and standard keys. The meristic and morphometric characters gathered fishes were estimated and checked and distinguished up to the species level.

The relative wealth was evaluated just for monetarily and economical essential fishes, which favored by customer. The overall bounty of individual species was determined by the accompanying equation:

$$\frac{\text{Number of sample of particular species}}{\text{Total Number of samples}} \times 100$$

Table 1: Canonical correlation matrix with two axes of the environmental variables in the canonical correspondence analysis (CCA) for three sites in the river Ganga

Axis variables	Correlation coefficient	
	1	2
Water Temperature(°C)	0.3391	0.9408*
pH	0.1879	-0.9822*
Total Dissolved Solid (mg ^l ⁻¹)	0.5334	-0.8459
SO ₄ (mg ^l ⁻¹)-1	-0.6112	-0.7914
PO ₄ (mg ^l ⁻¹)-1	0.9598*	-0.2806
Alkalinity (mg ^l ⁻¹)	-0.8865	0.4628
Total Hardness (mg ^l ⁻¹)	-0.9966*	0.0828
Nitrate (mg ^l ⁻¹)	0.9799*	-0.1994
Dissolved Oxygen (mg ^l ⁻¹)	-0.3241	0.9460*
Biological Oxygen Demand (mg ^l ⁻¹)	-0.8142	-0.5806

Cadmium (mg ^l ⁻¹)	-0.6368	-0.7710
Zink (mg ^l ⁻¹)	-0.4086	-0.9127*
Lead (mg ^l ⁻¹)	-0.8182	-0.5749




Figure 1: Canonical Correspondence Analysis (CCA). The multivariate analysis indicates the relationship between fish species and environmental variables in the River Ganga. Fish species and variables are indicated by filled circles and arrows, respectively

4. Result and Discussion

4.1 Canonical correspondence analysis (CCA)

We break down information in each of the three destinations all in all stretch since we evaluated a go along results for all locales. A CCA chart does not have to contain every one of the components (animal varieties, locales, environmental factors). To abstain from congestion of focuses, species and locales are regularly appeared separate outlines that can, on a basic level, be overlain. On the other hand, chose focuses or factors are shown. Sanctioned correspondence analysis (CCA) demonstrated that hub 1 and 2 represented 67% and 33% difference for species and environmental connection, individually. The biplots measurements created for every one of the three stations by CCA recommended that all out hardness was most imperative factor at pivot 1, while Nitrate and Phosphate were additionally essential at a similar hub. At pivot 2, pH was most critical factor pursued by broke down oxygen; water temperature and Zink metal (Table 1). These factors were connected altogether (p=0.6660, F-value=0.67) for hub 1 and 2. Complete hardness, alkalinity and broke up oxygen were in charge of the nearness of catla, rita and Sperataaor, while Labeocalbasu, Cyprinus carpio and Cirrhinus mrigala favored nitrate, phosphate and all out disintegrated strong for their bounty. Oreochromis niloticus favored high organic oxygen request and lead while Zn and Sulfate were in charge of wealth of L. rohita.

Appointment analysis uncovered that environmental factors impact substantially the fish fauna in the Ganga stream; all out hardness, nitrate, phosphate, DO, pH and water temperature and Zn metal were most imperative factors for the bounty of L. rohita, L. calbasu, C. catla, C. mrigala, R. rita and C. carpio. O. niloticus was overwhelming metal favored fish. Environmental conditions impact fish appropriations, networks and occasional developments. To limit vitality exhausted for

survival, species commonly support territories that advance their physiological procedures. Moyle et al., Bain et al., Lobb et al. likewise revealed water profundity, momentum speed and substratum as essential factor for the plenitude of R. alburnoides and L. pyrenaicus in the American streams.

4.2 Structure of the fish array

Fish arrays in the Ganga stream arrange are affected by both confined territories and bigger scene examples and water management system. Real neighborhood factors are 1) accessibility of various kinds of natural surroundings condition, 2) accessibility of various sorts of sustenance living beings and assets, and 3) communications with other amphibian species (e.g., predation, focused cooperations). About portion of Indian fishes are in the minnow family (Cyprinidae). Amid the investigation time frame diverse fish assortments have been recorded in the Ganga waterway at Kanpur, Allahabad and Varanasi destinations, India. Human exercises and mechanical influent most extreme announced in these destinations of the stream, so we have picked these locales. The outcome demonstrated that the region was wealthy in fish decent variety. Fish biodiversity of the Gang River from Kanpur to Varanasi harbors of 102 fish species (with assortment) have a place with 8 requests and 28 families (Table 2). Cypriniformes and Cyprinidae were the most rich species request and family. At complete stretch, Cypriniformes request was shared 49 species (48.04%), trailed by Siluriformes 26 species (25.49%) and Perciformes 17 species (16.67%). Requests Clupeiformes shared 5 species (4.90%) (Figure 2)

At Kanpur site, 74 fish species were recorded with 6 orders. Cypriniformes request was shared 32 species (43.24%) trailed by Siluriformes 21 species (28.38%) and Perciformes 15 species (20.27%). Request Clupeiformes shared 3 species (4.05%). At Allahabad site, 89 fish species were recorded with 8 orders. Cypriniformes request was shared 41 species

(46.07%) trailed by Siluriformes 25 species (28.09%) and Perciformes 14 species (15.73%). Request Clupeiformes and Osteoglossiformes shared 4 species (4.49%) and 2 species (2.25%), separately (Figure 4). At Varanasi site, 82 fish species were recorded with 7 orders. Cypriniformes request was shared 39 species (47.56%) trailed by Siluriformes 23 species (28.05%) and Perciformes 13 species (15.85%). Request Clupeiformes and Osteoglossiformes shared 3 species (3.66%) and 2 species (2.44%), individually (Figure 5). Cyprinus carpio var. communis and Oreochromis niloticus are much of the time recorded in the Ganga waterway. The two species had extensive dispersal limit. The two species are intriguing/outsider fish species for India. For protection perspective C. carpio var. communis and O. niloticus species ought to be checked in the Ganga River. The two species are extremely hurtful for fish biodiversity in any substantial water bodies as like waterways, lakes and repositories. Fishes are undermined by channelization of waterways/streams beds.

Table 2: Biodiversity of fishes from the Ganga River at Kanpur to Varanasi, India

S. N.	Order/Family/Genus/Species	Kanpur	Allahabad	Varanasi
	Order- Osteoglossiformes			
	Family: Notopteridae			
1	Chitalachitala	+	+	+
2	Notopterusnotopterus	+	+	+
	Order- Anguilliformes			
	Family: Anguillidae			
3	Anguilla bengalensis		+	
	Order- Clupeiformes			
	Family: Clupeidae			
4	Gudusiachapra	+	+	+
5	Goniolosamanmina	+	+	
6	Tenualosa (Hilsa) ilisha			+
	Family: Pristigasteridae			
7	Ilishamegaloptera		+	
	Family: Engraulidae			
8	Setipinnaphasa	+	+	+
	Order- Cypriniformes			
	Family: Cyprinidae			
9	Catlatcatla	+	+	+
10	Chaguniuschagunio	+	+	+
11	Cirrhinusmrigala	+	+	+
12	Cirrhinusreba	+	+	+
13	Ctenopharyngodonidella	+	+	+
14	Hypophthalmichthysmolitrix	+	+	+
15	Cyprinus carpio communis	+	+	+
16	Cyprinus carpio specularis	+	+	
17	Aristichthysnobilis	+		+
18	Labeoangra		+	+
19	Labeocalbasu	+	+	+
20	Labeobata	+	+	+
21	Labeoboga	+		

22	Labeorohita	+	+	+
23	Labeogonius		+	+
24	Labeopangusia		+	
25	Osteobramabelangeri			+
26	Osteobramacotiocotio	+	+	+
27	Puntiuschola		+	+
28	Puntiusconchoniunus			+
29	Puntius saranasarana	+	+	+
30	Puntius sophera	+	+	+
31	Puntius ticto	+	+	+
32	Chela laubuca	+	+	+
33	Chela sladonii		+	+
34	Chela cachius		+	
35	Salmostomabacaila	+	+	+
36	Salmophasiaphulo	+	+	
37	Amblypharyngodonmola	+	+	+
38	Aspidopariajaya	+	+	+
39	Aspidopariamorar	+	+	+
40	Bariliusbarila	+	+	+
41	Bariliusbarna		+	
42	Bariliusbendelisis		+	+
43	Bariliusbola	+	+	+
44	Bariliusvagra		+	
45	Esomus danricus			Yes
46	Rasborarasbora	+	+	+
47	Raiamasbola		+	+
48	Tor tor		+	
49	Securiculagora	+	+	+
50	Osteobramacotiocotio			+
51	Crossocheiluslatiuslatius	+	+	+
	Family: Balitoridae			
52	Nemacheilusbotia	+	+	+
53	Aborichthyselungatus		+	
	Family: Cobitidae			
54	Botiaalmorhae		+	+
55	Botialohachata	+	+	+
56	Botiadarior	+		+
57	Lepidocephalusguntea	+		
	Order- Siluriformes			
	Family: Bagridae			
58	Sperataaor	+	+	+
59	Sperataseenghala	+	+	+
60	Mystustengra	+	+	+
61	Mystuscavasius	+	+	+
62	Mystusvittatus	+	+	+
63	Mystusbleekeri		+	
64	Rita rita	+	+	+
	Family: Siluridae			
65	Ompokbimaculatus		+	+

66	Ompakpabda	+	+	+
67	Wallagoattu	+	+	+
	Family: Schilbeidae			
68	Ailiacoila	+	+	+
69	Clupisomagaru	+	+	+
70	Eutropiichthysvacha	+	+	+
71	Eutropiichthysmurius	+	+	+
72	Siloniasilondia	+	+	+
	Family: Pangasiidae			
73	Pangasiuspangasius		+	+
	Family: Sisoridae			
74	Bagariusbagarius	+	+	+
75	Gagatacenia	+	+	+
76	Nangranangra			
77	Nangraviridescens	+	+	+
78	Sisorrhodophorus	+	+	+
79	Glyptothoraxlineatus		+	
	Family: Clariidae			
80	Clariasbatrachus	+	+	+
81	Clariasgariepinus	+	+	+
	Family: Heteropneustidae			
82	Heteropneustesfossilis	+	+	+
	Family: Belonidae			
83	Xenentodoncancila	+	+	+
	Order-Synbranchiformes			
	Family: Synbranchidae			
84	Monopteruscuchia		+	+
	Order-Perciformes			
	Family: Ambassidae			
85	Chandanama	+	+	+
86	Chandaranga	+		+
87	Pseudambassisranga			+
	Family: Sciaenidae			
88	Johniuscoitor	+	+	+
89	Family: Mugilidae			
90	Rhinimugilcorsula	+	+	+
	Sicamugilcascasia	+	+	+
	Family: Gobiidae			
91	Glossogobiusgiuris	+	+	+
	Family: Anabaniitidae			
92	Anabas testudineus	+	+	+
	Family: Belontiidae			
93	Colisafasciatus	+	+	
	Family: Channidae			
94	Channamarulius	+	+	+
95	Channapunctatus	+	+	+
96	Channastratus	+	+	+
97	Channastewartii	+		
	Family: Mastacembelidae			

98	Macragnathuspancalus	+	+	
99	Mastacembelusarmatus	+	+	+
	Family: Nandidae			
100	Nandusnandus		+	
	Family: Cichlidae			
101	Oreochromisniloticus	+	+	+
	Order- Tetraodontiformes			
	Family: Tetraodontidae			
102	Tetraodoncutcutia	+	+	+
Total	102	74	89	82

More species are expected to protect a steady supply of ecosystem goods and services as spatial and temporal changeability expands, which commonly happens as longer timeframes and bigger regions are considered. Every natural environment has an assortment of animal categories, which contrast in their relative wealth. No people group comprises of types of equivalent wealth. A few animal categories are uncommon, others are normal and still others might be bounteous. Nautiyal et al. recorded 122 fish species from the Ganga River (Haridwar to Kanpur segment). Menon has recorded 207 types of fish from the Gangetic fields which have a place with 29 families and 82 genera. As per another gauge, the Gangetic system alone harbors at the very least 265 types of fish. Freshwater biodiversity has declined quicker than either earthy or marine biodiversity in the course of recent years. Presentations of non-indigenous fishes can diminish assorted variety and change nearby network elements in freshwater systems. The physical and organic attributes of riverine systems have been appeared to shape fish network.

4.3 Abundance of some important fishes

Abundance was recorded only commercially important fish species, which preferred by consumer and had high market price. Out of 102 species, species having higher economical are *C.catla*, *L.rohita*, *C. mrigala*, *L. calbasu*, *S. aor*, *S. seenghala*, *W. attu*, *R. rita*, *E. vacha*.




Figure 2: Contribution of different orders at Kanpur to Varanasi section




Figure 3: Contribution of different orders at Kanpur site




Figure 4: Contribution of different orders at Allahabad site




Figure 5: Contribution of different orders at Varanasi site

Table 3: Relative abundance of some commercially important fishes from the Ganga River

Fishes	Kanpur		Allahabad		Varanasi		Total	
	Occurrence	%	Occurrence	%	Occurrence	%	Total number	%
Major carp								
Catlacatla	106	2.32	187	2.15	96	0.89	389	1.62
Labeorohita	163	3.56	137	1.57	179	1.67	479	1.99
Cirrhinusmrigala	208	4.55	311	3.57	463	4.32	982	4.08
Labeocalbasu	123	2.70	196	2.35	407	3.80	726	3.02
Catfishes								
Sperataaor	463	10.12	960	11.01	1007	9.39	2430	10.11
Sperataseenghala	721	15.76	1323	15.17	1602	14.94	3646	15.18
Wallagoattu	226	4.94	401	4.60	521	4.86	1148	4.78
Rira rita	183	4.00	486	5.57	372	3.47	1041	4.33
Eutropiichthysvacha	638	13.95	1630	18.69	2164	20.19	4432	18.46
Clupisomagaru	671	14.67	1482	17.00	2063	19.25	4216	17.62
Exotic fishes								
Cyprinuscarpiovarcommunis	469	10.25	763	8.75	1082	10.09	2314	9.64
Oreochromisniloticus	603	13.18	843	9.67	763	7.12	2209	9.19
Total	4574		8719		10719		24012	

C. garua and C. carpio var. communis and O. niloticus had moderate economic esteem. C. catla, L. rohita, C. mrigala, L. calbasu are herbivorous in encouraging while S. aor, S. seenghala, W. attu, R. rita, E. vacha. C. garua are rapacious in sustaining. C. carpio var. communis and O. niloticus are omnivorous in bolstering.

Standard correspondence analysis affirmed measurably exceptionally noteworthy contrasts ($P < 0.0001$) between fish wealth of the individual site. Altogether, 21.2% of fish array inconstancy is clarified by this example, as it generally outlines contrasts in environmental states of individual site. C. carpio var. communis and O. niloticus the two species are intruder species in the Ganga River. As indicated by pooled bounty E.

vacha was ruled fishes in the complete stretch from the Ganga River. Indian real carp bounty was poor at present work. Catfishes were commanded in carp groups. Bounty of extraordinary species was likewise high. At Kanpur site, *S. seenghala* (15.76%) was ruled contrasted with *C. garua* (14.67%) and *E. vacha* (13.95%). Its seemed 2.32%, 3.56% and 4.55% of *C. catla*, *L. rohita* and *C. mrigala*, separately. *C. catla*, *L. rohita* and *C. mrigala* bounties were unequivocally associated with temperature. At Allahabad site, *E. vacha* (18.69%) was overwhelmed contrasted with *C. garua* (17.00%) and *S. seenghala* (15.17%). At Varanasi site, *E. vacha* (20.19%) was commanded contrasted with *C. garua* (19.09%) and *S. seenghala* (14.94%). Present examination demonstrated that the *O. niloticus* unequivocally connected with high organic oxygen request and lead. Lakra et al watched relative wealth 1.33, 2.75, 1.21 and 0.34 of *C. catla*, *L. rohita*, *C. mrigala* and *C. carpio* from the Betwa River. *O. niloticus* and *C. carpio* intensely attacked in the Ganga River and its biggest tributary the Yamuna River. Colorful species may become obtrusive and are equipped for spreading intriguing ailments, diminishing biodiversity through competition, predation and natural surroundings corruption, hereditary weakening of wild populaces through hybridization and quality introgression in short or long course of time.

5. Conclusion

Range expansion of specific species and decrease in scopes of couple of species is a genuine worry in the long haul preservation of fishes in the Ganges. In addition, higher wealth of exotics, fracture and changes in the hydrology of stream due to hydro activities and boundaries are real danger to the fishes in the Ganges separated from unpredictable fishing, contamination, poor land use design. Up until this point, in India fishes are considered as business item and diminished value of their ecological services which pushed expansive number of species under compromised classifications. Fish preservation zones, scene level protection plan, appropriate Environment Impact Assessment for any formative exercises in the bowl, natural surroundings rebuilding plan, species recovery plan for certain undermined species in the Ganges and so forth may help the local fish decent variety reestablish in the Ganga. The Landing information of the Ganga River likewise demonstrated that the ecological states of the stream are falling apart for fish. The waterway Ganga has been recently known as unique adobe of the important Indian real carps. The arrival of Indian real carps has fundamentally declined. The regular variety likewise impacts the arrival from the Ganga River.

References

- Allan JD, Abell R, Hogan Z, Revenga C, Taylor BW, et al. (2005) Overfishing of inland waters. *Bioscience* 55:1041-1051.
- Priyanka M, Dwivedi AC (2015) *Biology of Cirrhinus mrigala and Oreochromis niloticus*. LAP LAMBERT Academic Publishing, Germany.
- Leveque CT, Oberdorff D, Paugy MLJ, Tedesco PA (2008) Global diversity of fish (Pisces) in freshwater. *Hydrobiologia* 595:545-567.
- Duncan JR, Lockwood JL (2001) Extinction in a field of bullets: a search for causes in the decline of the world's freshwater fishes. *Biological Conservation* 102:97-105.
- Argent DG, Bishop JA, Stauffer JR, Carline RF, Myers WL (2003) Predicting freshwater fish distributions using landscape-level variables. *Fisheries Research* 60:17-32.
- Yakubu A, Li N, Conrad J, Zeeman ML (2011) Fish population dynamics and constant or periodic proportion harvesting policies. *Math Bioscience* 232: 66- 77.
- Lakra WS, Sarkar UK, Kumar RS, Pandey A, Dubey VK, et al. (2010) Fish diversity, habitat ecology and their conservation and management issues of a tropical River in Ganga basin, India. *Environmentalist* 30:306-319.
- Dwivedi AC, Nautiyal P (2012) Stock assessment of fish species *Labeo rohita*, *Tor tor* and *Labeo calbasu* in the Rivers of Vindhyan region, India. *Journal of Environmental Biology* 33:261-264.
- Junior L, Cardone IB, Goitein R (2006) Fish assemblage structure and aquatic pollution in a Brazilian stream: some limitations of diversity indices and models for environmental impact studies. *Ecology of Freshwater Fishes* 15:284-290.
- Pathak RK, Gopesh A, Joshi KD (2013) Alien fish species, *Cyprinus carpio* var. *communis* (common carp) as a powerful invader in the Ganga River at Allahabad, India. *Journal of the Kalash Science* 1:133-139.
- Marti M, Berthou GE, Sabater S, Tomanova S, Monoz I (2010) Comparing fish assemblages and trophic ecology of permanent and intermittent researches in a Mediterranean stream. *Hydrobiologia* 215:167-180.
- Pathak RK, Gopesh A, Dwivedi AC (2011) Alien fish species, *Cyprinus carpio* var. *communis* (common carp) as a powerful invader in the Yamuna River at Allahabad, India. *National Academy of Science Letter* 34:367-373.
- Sinha M, Khan MA (2001) Impact of environmental aberrations on fisheries of the Ganga (Ganges) River. *Aquatic Ecosystem Health and Management Society* 4:493-504.
- Mantana CG, Choudhary SK, Dey S, Winemiller KO (2011) Compositional trends of fisheries in the River Ganges, India. *Fisheries Management and Ecology* 18:282-296.
- Nautiyal P, Mishra AS, Singh KR, Singh U (2013) Longitudinal distribution of the fish fauna in the River Ganga from Gangotri to Kanpur. *Journal Applied National Science* 5:63-68.
- Lobb MD, Orth DJ (1991) Habitat use by an assemblage of fish in a large warm water stream. *Transactions of the American Fisheries Society* 120:65-78.
- Dwivedi AC, Tewari NP, Mayank P (2007) Biodiversity of fishes of Faizabad District. *Flora and Fauna* 13:379-383.
- UNESCO (2003) *Water for People, Water for life*. The United Nations World Water Development Report, Paris.
- Dwivedi AC, Tiwari A, Mayank P (2015) Seasonal determination of heavy metals in muscle, gill and liver tissues of Nile tilapia, *Oreochromis niloticus* (Linnaeus, 1758) from the tributary of the Ganga River, India. *Zoology and Ecology* 25:166-171.
- Casal CMV (2006) Global documentation food fish introductions: the growing crisis and recommendations for action. *Biological Invasions* 8:3-11.
- Singh AK, Lakra WR (2006) Alien fish species in India: Impact and emerging