

# Study on Heterocyclic Compounds as Drugs and General Procedure for the Synthesis Of 4, 6- Disubstituted-1, 2, 4-Triazolo-1,3,4-Thiadiazoles

<sup>1</sup>Irudayaraj & <sup>2</sup>Dr. Komal Lata Nagpal

<sup>1</sup>Research Scholar, OPJS University, Churu Rajasthan (India)

<sup>2</sup>Assistant Professor, OPJS University, Churu Rajasthan (India)

## ARTICLE DETAILS

### Article History

Published Online: 12 June 2019

### Keywords

Thiadiazoles, medications, technique, antidepressants, heterocyclic.

## ABSTRACT

Heterocyclic chemistry is a significant part of organic chemistry representing about 33% of current distributions. Truth be told 66% of organic compounds are heterocyclic compounds. Thiadiazoles are a significant class of heterocyclic compounds that show various applications in pharmaceutical and natural applications. In this paper we will learn about the heterocyclic compounds as medications and general technique for the union of 4, 6-disubstituted-1, 2, 4-triazolo-1,3,4-thiadiazoles.


## 1. Introduction

Heterocyclic compounds are outstanding for their organic exercises. The primary source of these heterocyclic compounds is plants. Wide scope of heterocyclic compounds, utilized against different diseases, has their application in pharmaceutical businesses. The immense interest of such compounds can't be satisfy from normal source plants. Subsequently they are orchestrated utilizing different

courses and utilized. Heterocycles structure by a wide margin the biggest and old style divisions of organic chemistry.


### 1.1 Heterocyclic compounds effectively utilized as drugs

Sumatriptan, a heterocyclic compound is the first antimigrain medicate, substitution of sulfonamide moiety in sumatriptan with 1, 2,4-triazole, which is additionally an intense 5-HT<sub>1D</sub> receptor agonist.




### 1.2 Heterocyclic compounds as antidepressants

An antidepressant is a mental drug used to ease disposition issue, for example, real discouragement, and dysthymia. Medications including the monoamine oxidase inhibitors (MAOIS), tricyclic antidepressants (TCAs), tetracyclic antidepressants (TeCAs), particular serotonin reuptake inhibitors. Albeit for the most part less adequate than the tricyclic (heterocyclic) antidepressants in the treatment of torment, the SSRIs have a demonstrated viability in this clinical setting. Their absence of reactions in respect to the TCAs settles on the SSRIs a decent decision for those torment patients who can't or won't endure the symptoms of the TCAs, yet at more prominent money related expense.





### (a) Heterocyclic compounds as anthelmintic

Helminthes are parasitic worms, which taint an expected two billion individuals around the world, almost all in poor creating tropical or semitropical nations. Helminthic infections add to unhealthiness, pallor, hindered growth, intellectual debilitation, and expanded defenselessness to different diseases.



### (b) Heterocyclic compounds as antiulcer agents


Pyridine ring assumes a significant job in human digestion because of its collaboration with amino acids. A large number of the dynamic medications in the market contain pyridine moiety. Benzimidazoles likewise are notable for their pharmacological properties



### (c) Antibiotics

"Antibiotics" originates from the Greek enemy of ("against") and profiles ("life"). Antibiotics are drugs that either decimate bacteria or forestall their generation. Antibiotics that eliminate bacteria are designated "bactericidal" and the ones that stop the growth of bacteria are classified "bacteriostatic". Antibacterial antibiotics can be sorted dependent on their objective explicitness: "slender range" antibiotics target specific kinds of bacteria, for example, Gram-negative or


Gram-positive bacteria, while expansive range antibiotics influence a wide scope of bacteria. Antibiotics which focus on the bacterial cell divider (penicillins, cephalosporins), or cell film (polymyxins), or meddle with fundamental bacterial proteins (quinolones, sulfonamides) for the most part are bactericidal in nature. Those which target protein amalgamation, for example, the aminoglycosides, macrolides and antibiotic medications are typically bacteriostatic.



### (d) Antihypertensive Drugs

The control of blood pressure requires a consistent change of cardiovascular yield and fringe vascular obstruction. Antihypertensive Drugs are a class of drugs that are utilized in prescription and pharmacology to treat hypertension.

Chlorothiazide (I), flumethiazide (II), trichloromethiazide (III) and polythiazide (IV) are diuretic medication of the thiazide class that demonstrations by repressing the kidneys' capacity to hold water. This lessens the volume of the blood, diminishing blood come back to the heart and along these lines cardiovascular yield is accepted to lower peripheral.



I- R = -CHCl<sub>2</sub>, R<sub>1</sub> = -H II- R = -CH<sub>2</sub>SCH<sub>2</sub>CF<sub>3</sub> : R<sub>1</sub> = -CH<sub>3</sub> III-R=Cl IV-R=CF<sub>3</sub>

## 2. Thiadiazole Derivatives

Thiadiazoles are significant class of heterocyclic compounds that show different application in organic synthesis, pharmaceutical and natural applications. Different organic exercises of thiadiazole and their subordinates have been known from mid 1950s and from that point forward. The examination chip away at this heterocyclic system has prompted critical advancements in their chemistry and

science. Thiadiazole and their subordinates can be considered as straightforward five membered heterocycles having one sulfur and two nitrogen atoms. They are outstanding as metal chelating agents and hostile to consumption agents.

Thiadiazoles are a significant class of heterocyclic compounds that display different applications in pharmaceutical and organic applications. Because of their

wide range of utilizations researcher over the globe are taking a shot at this moiety. Subordinates of 1,3,4-thiadiazole and 1,2,4-triazole consolidated core system found to have assorted natural exercises, for example, insecticidal, bactericidal, herbicidal, calming, antiviral, fungicidal and CNS stimulant properties. As of now,

heterocycles bearing a 1,2,4-triazole or 1,3,4-thiadiazole moiety, speak to an intriguing class of compounds having a wide range of pharmacological exercises viz., mitigating, antiviral and antimicrobial. In addition, broad survey of writing has uncovered that s-triazolo [3,4-b]-1,3,4-thiadiazole rings, have gotten much consideration during ongoing years by virtue of their unmistakable usage as antifungal, calming, antiviral, pain relieving, antihelminthic and antibacterial agents. A triazolothiadiazole system containing – N C S linkage may go about as a cyclic simple of two significant parts—thiosemicarbazide and biguanide which regularly show differing organic exercises. In the present examination, we report the synthesis of a novel series of 4, 6-substituted 1, 2, 4-triazolo-1, 3, 4-thiadiazole subordinates. Also, we found that one of the compound 5-(3-(2,3-dichlorophenyl)-[1,2,4]triazolo[3,4-b][1,3,4]thiadiazol-6-yl)-2-fluorobenzonitrile (3f) displayed huge antitumor impacts by improving TNF-induced apoptosis through the negative guideline of NF- $\kappa$ B flagging course.

### 3. Materials and methods


actin immune response and paclitaxel were procured MTT, Tris, glycine, SDS, BSA, from Sigma-Aldrich (St. Louis, MO). Thiadiazole subsidiaries were broken down in dimethyl sulfoxide (DMSO) as a 20 mM stock arrangement and put away at 4°C. Most extreme last grouping of DMSO was under 0.1 % and utilized for control tests. Further weakening was done in cell culture medium as required. Dulbecco's adjusted Eagle medium (DMEM), fetal cow-like

serum (FBS) and anti-toxin antimycotic blend were acquired from Life Technologies.


Silica gel GF-254 was utilized for dainty layer chromatography.  $^1\text{H}$  range was recorded at 400MHz and Melting focuses were resolved utilizing SELACO-650 hot stage dissolving point mechanical assembly and were uncorrected. Turn multiplets are given as s (singlet), d (doublet), t (triplet) and m (multiplet). Mass and immaculateness were recorded on a LC-MSDTrap-XCT. Silica gel section chromatography was performed utilizing Merck 7734 silica gel (60–120 work) and Merck made TLC plates. The yields were disconnected yields and are the normal of two runs. All separated solvents were dried over  $\text{Na}_2\text{SO}_4$  and dissipated with a BUCHI rotating evaporator. Reagents were acquired economically and utilized as got.

#### 3.1 General procedure for the synthesis of 4, 6-disubstituted-1, 2, 4-triazolo1,3,4-thiadiazoles: 3(a-m)

We have synthesized the compound, 4-amino-5-(2, 3-dichlorophenyl)- 4H-1, 2, 4-triazole-3-thiol (1) by utilizing a revealed procedure George T (Scheme 2.2). The second arrangement of synthesis includes, a blend of 4-amino-5-(2, 3-dichlorophenyl)- 4H-1, 2, 4-triazole-3-thiol (1) and various acids 2(a-m) and  $\text{POCl}_3$  (5 mL) were refluxed medium-term. After finish of the response (hexane: ethyl acetic acid derivation: 7.5:2.5 utilized for TLC), the response blend was cooled and emptied gradually with mixing into super cold water. The pH of the mass was acclimated to essential (pH = 7.5-8.0) with  $\text{NH}_4\text{OH}$  arrangement and the subsequent compound was extricated with ethyl acetic acid derivation. The organic layer was washed with 10 %  $\text{NaHCO}_3$  answer for evacuate the un-responded acid, dried with  $\text{Na}_2\text{SO}_4$ , sifted and the dissolvable was dissipated under vacuum. To the buildup got, 3 volumes of hexane were included, mixed well and separated at virus condition.



Scheme 1.1: Possible mechanism for the synthesis of thiadiazoles 3(a-m)



#### 4. Results

In the present work, a series of 1,3,4 thiazadiazole subordinates 3(a-m) were synthesized by Scheme 2.2 (Scheme 2.1 clarifies the conceivable component for the synthesis of thiazadiazoles 3(a-m)).

#### 5. Conclusion

thiazadiazole subordinate can be a potential candidate in treating cervical cancer explicitly. Furthermore research is required to warrant the equivalent. It would likewise be

fascinating to survey the capability of 3f and related compounds in instigating apoptosis among assortment of cancer cell lines to advance cancer therapy utilizing thiazadiazole subordinates. Most of pharmaceuticals and naturally dynamic agrochemicals are heterocyclic while innumerable added substances and modifiers utilized in industrial applications extending from beauty care products, reprography, information stockpiling and plastics are heterocyclic in nature.

#### References

- [1] Nagender, P.; Malla Reddy, G.; Naresh Kumar, R.; Poornachandra, Y.; Ganesh Kumar, C.; Narsaiah, B. *Bioorganic & Medicinal Chemistry Letters* 2014, 24, 2905.
- [2] Sashidhara, K. V.; Modukuri, R. K.; Choudhary, D.; Bhaskara Rao, K.; Kumar, M.; Khedgikar, V.; Trivedi, R. *European Journal of Medicinal Chemistry* 2013, 70, 802.
- [3] El-Sayed, N. S.; Shirazi, A. N.; El-Meligy, M. G.; El-Ziaty, A. K.; Rowley, D.; Sun, J.; Nagib, Z. A.; Parang, K. *Tetrahedron Letters* 2014, 55, 1154.
- [4] Conti, C.; Proietti Monaco, L.; Desideri, N. *Bioorganic & Medicinal Chemistry* 2014, 22, 1201.
- [5] Moraski, G. C.; Oliver, A. G.; Markley, L. D.; Cho, S.; Franzblau, S. G.; Miller, M. J. *Bioorganic & Medicinal Chemistry Letters* 2014, 24, 3493.
- [6] Qin, M.; Zhai, X.; Xie, H.; Ma, J.; Lu, K.; Wang, Y.; Wang, L.; Gu, Y.; Gong, P. *European Journal of Medicinal Chemistry* 2014, 81, 47.
- [7] J. Ferlay, I. Soerjomataram, R. Dikshit, S. Eser, C. Mathers, M. Rebelo, D. M. Parkin, D. Forman and F. Bray, *International journal of cancer. Journal international du cancer*, 2015, 136, E359-386.
- [8] A. Jemal, F. Bray, M. M. Center, J. Ferlay, E. Ward and D. Forman, *CA: a cancer journal for clinicians*, 2011, 61, 69-90.
- [9] B.-R. L. Bruni L, Albero G, Aldea M, Serrano B, Valencia S, Brotons M, Mena M, Cosano R, Muñoz J, Bosch FX, de Sanjosé S, Castellsagué X, *Human Papillomavirus and Related Diseases in the World*. .
- [10] R. Sankaranarayanan, R. Swaminathan, H. Brenner, K. Chen, K. S. Chia, J. G. Chen, S. C. Law, Y. O. Ahn, Y. B. Xiang, B. B. Yeole, H. R. Shin, V. Shanta, Z. H. Woo, N. Martin, Y. Sumitsawan, H. Sriplung, A. O. Barboza, S. Eser, B. M. Nene, K. Suwanrungruang, P. Jayalekshmi, R. Dikshit, H. Wabinga, D. B. Esteban, A. Laudico, Y. Bhurgri, E. Bah and N. Al-Hamdan, *The Lancet. Oncology*, 2010, 11, 165-173.