

Study of Climatic Change Response of Egg Laying Behavior of *Antheraea Mylitta* D

Kumari Shobha Rani

Assistant Professor of Zoology, D N College of Masaurhi Patna

ARTICLE DETAILS

Article History

Published Online: 15 April 2019

Keywords

Traits, Humidity, Biological, Climatic, Range.

ABSTRACT

Antheraea mylitta Drury is a holometabolous bug with eggs, hatchling, pupa, and cover and grown-up are stages throughout its life cycle. High temperature above 28°C and less relative moistness than half are not appropriate for the oviposition and hatching of eggs. In Tasar silk moth, egg retention was likewise more even after 72 h of oviposition in normal conditions. From the investigations it is seen that rich biological diversity of *Antheraea mylitta* predominantly is because of its wide scope of conveyance, climatic factors, and food plants and so on., which have prompted made variations in their ethology, physiology and commercial qualities.

1. Introduction

Antheraea mylitta Drury, Andhra neighborhood ecorace is an exclusive race of Andhra Pradesh. Prior certain researchers have announced restricted positive reaction of *A. mylitta* for indoor rearing. Anyway more endeavors are to be made toward this path to rise from chawki stage to the casing stage. The accessible writing shows that Thangavelu et al. (1992) have revealed indoor chawki worm rearing of *A. mylitta* with an expected ERR (Effective pace of rearing) of fewer than 20%. In spite of the fact that the discoveries are not empowering, one angle evidently turns out to be certain that *A. mylitta* isn't unwilling to indoor rearing. Along these lines, an endeavor has been made to raise the bug from first instar to the last instar and allow it to go for case turning too. Present report gives subtleties of the moth weight concerning outside and indoor rearing.

Because of global warming, effect of temperature on living organisms has been center of logical examination in ongoing past. In creepy crawly, all species has its own selection of temperatures for its typical development and extremely high temperature hinders the development, may prompts developmental glitch, for example, larval ecdysis and grown-up rise. Being a poikilothermic organism, temperature chooses the destiny of creepy crawly, during embryonic and postembryonic development. It is surely known that, temperature influences the creepy crawly life-cycle pliancy, physiological thermal traits, immune reactions and quality articulation. It additionally influences the metabolic rate and oxidative pressure status considered the effect of temperature, water and oxygen on creepy crawly eggs at a progress among dissemination and reaction limitation. Indeed, even oxygen utilization profiles of bug cells get influenced because of temperature. In various bug species, the development rate is improved and the developmental period and immune cells (hemocytes) are compressed as the temperature rises.

The tropical tasar silkworm, *A. mylitta* is produces tasar silk of commercial significance at industry level and it gives direct advantageous occupation to needy individuals of the society. The essential regenerative unit in the tasar silk industry is the silkworm eggs and temperature assumes urgent job during embryonic development. Embryonic development begins inside scarcely any long stretches of egg lying and it needs legitimate hatching and care. It announced that, as the

development of the undeveloped organism continues, the two lipids and starches gets used, bringing about increment in the concentration of citrate, pyruvate and lactate. Because of global warming rise in temperature with less humidity was recorded during first harvest grainage which influence the egg hatching as well as unfavorably influences the seed division of tasar industry which is extremely fundamental contribution of sericulture. In spite of the fact that, embryology of a few sericigenous creepy crawlies have been concentrated by specialists. Yet, biochemical and morphological changes happens in tasar silkworm eggs under temperature stress conditions are not seen well. Subsequently, the current examination has been directed to investigate effect of temperature weight on embryonic development and biochemical profile of the *A. mylitta* eggs.

2. Literature Review

Satisfaction Kishan Sharma H (2017) The targets of this investigation on the "Biochemical examination of haemolymph of Tasar silkworms" is to look through standard available resources for improving silk execution traits which are required in generating rustic business work and earning of unfamiliar incomes. *Antheraea mylitta* Drury is a holometabolous bug with eggs, hatchling, pupa, case and grown-up are stages throughout its life cycle. There are five larval instars which feed on the essential host plants like Arjun (*Terminalia arjuna*), Asan (*Terminalia tomentosa*), Sal (*Shorea robusta*) separated with various host plants of optional significance. The hatchlings unquenchably feed on the leaves of their host plants and fifth instars hatchling turns the silk around it and structures the casing from which Tasar silk is reeled. Silkworms are being utilized as bio-manufacturing plant for the production of valuable silk proteins which are normal polymers and are biodegradable, with receptive practical gatherings that open possibilities to connect the yields with different polymers to be utilized in controlled delivery framework.

Madhusudhan KN (2017) Due to open air rearing of tasar silkworm, the hatchlings are amiable to various abiotic and biotic pressure conditions which lead to extreme mortality of the hatchlings. Three distinctive tasar silkworm rearing fields were chosen for the current examination. The a-biotic factors viz., Temperature, Relative humidity and precipitation were recorded normally alongside dead hatchlings because of

various illnesses and irritations. The information gathered were completely investigated. The outcomes uncovered that, vacillation in nature prompts expanded sickness occurrence. The vermin pervasion was extreme during the early instars of the silkworm. Existing field conditions alongside fluctuating abiotic factors assume significant job in the survivability of tasar silkworm hatchlings in open air condition.

Renuka. G (2013) The polyphagous sericigenous bug, tasar silkworm, *Antheraea mylitta* Drury is generally found in the tropical regions of our nation, taking care of principally on food plants like *Terminalia arjuna* and *T. tomentosa* a variety of optional food plants. Tasar culture is a conventional livelihood for lakhs of tribal populace in the territories of Jharkhand, Chhatisgarh, Orissa, Maharashtra, Andhra Pradesh, West Bengal and Uttar Pradesh. The current investigation depicts the variation in the various phases of tasar silkworm *Antheraea mylitta* Drury which is conveyed as ecoraces in fluctuated geographical territories.

P. Sudhakara Rao (2019) Tasar silkworm (*Antheraea mylitta* Drury) rearing and casing production are basically completed by Tribal individuals of Central India and a few pieces of Odisha and Telangana states. Absence of Tasar silkworm seed production and conservation innovation is making difficulties in opportune gracefully of satisfactory quantity of seed to the ranchers and others stake holders..High temperature above 28°C and less relative humidity than half are not appropriate for the oviposition and hatching of eggs. In Tasar silk moth, egg retention was additionally more even after 72 h of oviposition in common conditions. In the current examination, while giving ideal temperature (27-28°C) and humidity (75-80%) in the egg laying room, egg lying was completed in bivoltine and trivoltine agonizes under various pressure conditions. The outcomes demonstrated that in bivoltines (per5 moths), in T1 where antenna and wings were excised, the quantity of eggs recorded were 1306, in T2 (legs and wings cut) it were 902 and in T3 (antenna, wings and legs cut), the quantity of laid eggs were 1154 when contrasted with 907 in Normal conditions (T0) for 5 months. Egg retention results demonstrated that in T1, it was 200 adding up to 13.28% and fruitfulness was 261. In T2, it was 145 adding up to 13.85 % and fertility was 180. In T3, with Antenna, legs and wings removal egg retention was 135 adding up to 10.47 % and fertility of 231. In instance of typical moths (T0), egg retention was 124 adding up to 12.03 % with a fruitfulness of 181. In the event of Trivoltines (5 moths) likewise same example was watched. The referenced pressure conditions are helpful while grainage tasks for getting greater fertility.

K. B. SHARMA (2013) Laboratory culture of Indian tasar silkworm, *Antheraea mylitta* has uncovered that the environmental factors viz; quality of food, Photoperiod, temperature, relative humidity and furthermore the period of rearing have significant effect on the social exercises of tasar silkworm. Consequences of the investigation have indicated that the develop quality of leaf, 16h. Photoperiod, 30°C temperature, 80% relative humidity and commercial yield season are a portion of the ideal conditions required for the healthy development and development of tasar silkworm under Indoor condition. The aggregate impacts of previously mentioned appropriate factors in mix have been found highly huge in relation to rearing exhibitions of tasar silkworm. Apparently the essential environmental factors are expected to

give required helpful and ideal condition for the best possible modification and acclimatization of tasar silkworm under indoor condition. Anyway tasar moths have been found to favor brief day photoperiod for their reproducing exercises.

3. Effect of rearing seasons on biological and economical traits

Season demonstrates the between yearly variation in temperature, humidity, daylight, rainfall, and so forth of a specific spot, which is represented by various geographical boundaries. Studies have demonstrated the significance of seasonal variations in science and development of creepy crawly. Bugs are merciless (Poikilothermic) organisms, so their internal heat level is pretty much same as that of their encompassing microenvironment. Temperature impacts all that an organism does, humidity influences embryonic development, and rainfall influences both. Environmental temperature, being the major abiotic factor, manages the internal heat level of caterpillar that decides the paces of taking care of, fertility, and mortality. Temperature is likely the absolute most significant environmental factor that impacts conduct, development, endurance, reproduction, and geographical dispersion of creepy crawlies. All the creepy crawly species has their own selection of temperatures for regular development and high temperature hinders the development that may prompts developmental breakdown, for example, larval ecdysis and grown-up emergence. Temperature systems and various degrees of relative humidity are known to assume a significant job in the existence pattern of bug and its versatility to the neighborhood atmosphere. A few scientists accept that the impact of temperature on bugs to a great extent overpowers the impacts of other environmental factors and that is the reason, the greater part of the creepy crawly's cautioning and anticipating models depend on collected degree-days from a base temperature and bio-fix point. Yamamura and Kiritani (1998) assessed that with a 2o C ascend in temperature, creepy crawlies may encounter one to five extra life cycles for every season. Most specialists appear to concur that hotter temperatures bring about more sorts and higher populaces of bugs. It has been accounted for that the temperature has highly critical impacts on all developmental boundary of bug than humidity and rainfall. Developmental season of a bug is contrarily identified with temperature.

Second abiotic factor that has huge effect on the presentation of creepy crawlies in earthly conditions is humidity. Humidity collaborates with free water accessibility and with the water substance of the food plants. There is no restricting scope of humidity and most bug can create at any humidity gave they can control their water balance. Humidity shows generally backhanded impact on development and development. Under too dry conditions, the leaves shrink extremely quick and become inadmissible for bug feed, bringing about hindered development of hatchlings, which makes them feeble and effectively helpless to sicknesses and other antagonistic conditions. During egg brooding, it is significant that humidity ought to be kept up at 80% on a normal for ordinary development of undeveloped organism. On the off chance that humidity falls underneath 70% during brooding, it influences hatching. Studies have likewise shown the joined impacts of temperature and relative humidity. Rainfall changes the working of microhabitat, which alongside

soil and other environmental factors influences foliage and water levels, which thusly influence on execution of bug. Artificially the variation from delicious leaves to develop is enormous to such an extent that it might divide the development pace of Lepidopteran hatchlings.

Change in temperature, humidity and other seasonal factors keeps creepy crawlies from achieving their physiological potential life history execution. Nonetheless, timberland bugs show a wonderful scope of transformations to changing situations and keep up their inner temperature (thermoregulation) and water content inside average cutoff points, regardless of wide vacillations in their environmental factors, since effect of temperature is altered by living space and other states of being. Temperature, humidity, air flow, gases and photograph period and so forth shows a noteworthy association in their impact on the physiology of silkworm relying on the mix of factors and developmental stage influencing development, development, profitability and quality of silk. Seasonal and environmental association have incredible task to carry out on the declaration of commercial characters of *A. mylitta*.

4. Materials & Methods

The current examination was embraced at the Breeding and Genetics Grainage house for the readiness of proportional crosses and rearing of tasar silkworm was led at all around kept up host plant field lab of Central Tasar Research and Training Institute, Ranchi, during second harvest (September-October) season.

Collection of cocoons for breeding: Daba semidomesticated cases were gathered after first yield reaping of Silkworm rearing and Genetics segment's germplasm bank, CTR&TI, Laria and Daba wild covers were gathered from its normal eco-specialties ie., Sal backwoods of Peterbar zone of Jharkhand and woods region close Hatgamaria, Jharkhand individually. All the casings were safeguarded independently in isolated out-entryway confines indicated for conservation of covers at the Grainage territory till the moth emergence.

Moth Emergence pattern: Moth emergence was concentrated during second grainage season (September), the quantity of moth developed and male - female synchronization was recorded in Daba (wild), Daba (S-d) and Laria ecoraces.

Coupling Behaviour: Coupling conduct of various crosses arranged was recorded dependent on oneself (regular) and hand coupling (initiated) of 50 every individual crosses in all the clumps. The coupling of moths was spoken to in the level of coupling.

Fecundity Assessment: Male and female mated moths of various blends were allowed for lovemaking for around 6-7 hours in standard grainage conditions. Female moths were decoupled and kept in paper box for ovi-position up to 72 hrs. The quantity of eggs laid by singular mother moth in three days (72 hrs.) was considered and recorded fertility. Systematic mother moth assessment was directed to identify any pebrine sickness. The eggs were washed to eliminate muconium layer and surface cleaned to plan ailment free layings.

Statistical analysis: Single direction ANOVA utilizing PC application SPSS 10.0 factual bundle to test the noteworthy of contrasts, between the methods for people and between the gatherings examined. Examinations were performed with Duncan's Multiple Range Test (DMRT, P< 0.05).

5. Results

The normal load of five male moths of outside reared *A.mylitta* (Andhra nearby) in the three harvests was 1.45, 1.49 and 1.42 gms, while that of females were 2.15, 1.99 and 2.18 gm. The normal load of male moths of indoor reared hatchlings in the three harvests was 1.44, 1.62 and 1.29 gm, though that of females was 2.08, 2.05 and 2.09. It is observed from the outcomes, that, male and female moth loads of outside reared hatchlings were slightly more prominent that those of indoor reared ones with the exception of in the subsequent yield (Table 1).

Table 1: Average moth weight of Tasar silkworm *Antheraea mylitta* (Andhra local) during three crops (in grams)

Crop	Sex	Outdoor rearing	Indoor rearing
I	Male	1.45	1.44
	Female	2.15	2.08
II	Male	1.49	1.62
	Female	1.99	2.05
III	Male	1.42	1.29
	Female	2.18	2.09

Table 2: Pupal and moth duration of irradiated cocoons of tasar silkworm *Antheraea mylitta* (Andhra local) reared under outdoor conditions

S.No	Dosage	Sex	Pupal duration after treatment (days)	Moth duration of life (days)
1	5 Gy (500 RADS)	Female	30	3
		Male	28	16
2	7.5 Gy (750 RADS)	Female	14	5
		Male	11	8
3	10 Gy (1000 RADS)	Female	12	6
		Male	10	10

In the moths of cases light at 5 Gy (500 RADS), the size of the wings and the body were littler than that of ordinary ones. The wing size had continuously diminished from first classification to third classification of radiation. In the moths of first classification, both the wings were collapsed, though in the subsequent kind, just the left wing was collapsed while in the third sort, the wings were collapsed, yet they were twisted and slender fit as a fiddle. The wings have demonstrated high beating recurrence and there were lost scales. The shade of wings of first and second classification moths was typical while that of third class was blurred and dull.

Cross-reproducing connotes a proficient technique for creature populace improvement. For silkworms, the cross-rearing is expected on one hand for the improvement of the varieties and then again for guaranteed affordable purposes comprising in delivering commercial crossovers, for which the prevalence against unadulterated varieties. The fruitfulness and egg ripeness are considered as one of the most wanted quantitative characters of commercial significance in silkworms. The genotype-condition connection related with host plant quality has highly huge influence on the fertility of the silkworms. The healthy and power of the mating male-female moths are significant for the subsequent quality and quantity of egg lying. Likewise the males of various varieties can influence the egg laying performance in silkworms. The fruitfulness of various equal crosses recorded in the current examination shows the fluctuation in the quantity of eggs laid by mother moths. Higher fruitfulness recorded in a portion of the crosses contrasted with the Daba (control). The impact of hybridization is a lot of evident and the due to the between ecorace cross. Additionally the heartiness and vigority of the wild relatives are influenced the higher fertility and quality of the eggs in the progeny.

In the current investigation the egg laying conduct and fruitfulness was observed in bivoltine and Trivoltine broods of

Daba race for 5 gravid female moths in every treatment under various pressure conditions. The outcomes demonstrated that for bivoltine (Table.1) with Antenna& wings removal (T1) recorded 766 eggs on first day and 540 on second day adding up to adding up to 86.72%. Egg retention for 5 months was 200 adding up to 13.28% and fruitfulness was 261 eggs for each month. In T2 with legs and wings cut away demonstrated 539 eggs on first day and 363 on second day adding up to and adding up to 86.15 %. Egg retention was 145 adding up to 13.85 % and fertility was 180. In T3, with Antenna, legs and wings cut demonstrated 1015 eggs on first day and 139 on second day adding up to and adding up to 89.53 % and egg retention was 135 adding up to 10.47 % and fertility of 231. In instance of typical moths (T0), it was shown 682 eggs on first day and 225 on second day adding up to, adding up to 87.97 % and eggs retention was 124 adding up to 12.03 % per with a fruitfulness of 213.

The outcomes showed for Trivoltine brood with Antenna& wings removal (T1) recorded 781 eggs on first day and 450 on second day adding up to summing to 95.42%. Egg retention was 59 adding up to 4.57% and fruitfulness was 246 eggs. In T2 with legs and wings severed showed 516 eggs on first day and 316 on second day adding up to and adding up to 85.15 %. Egg retention was 245 adding up to 25.07% and fertility was 166 every month. In T3, with Antenna, legs and wings cut demonstrated 833 eggs on first day and 293 on second day adding up to and adding up to 96.98% and egg retention was 35 adding up to 3.01 % and fruitfulness of 225. In instance of ordinary moths (T0), it was shown 582 eggs on first day and 328 on second day adding up to, adding up to 86.34 % and eggs retention was 144 adding up to 13.66 % with a fertility of 182. The load of male, female moths, single egg weight and eggs/g were recorded and delineated in Table 3 showing 95 eggs/gm and 85 eggs/gm in bivoltine and trivoltine separately.

Table.3 Egg laying efficiency in Bivoltine Tasar silk moths (sample 5 moths)

Treatments	1 st day	2 nd day	Total laid eggs		Egg retention in the abdomen		Total eggs	Fecundity (No)
			No	(%)	No	(%)		
Antenna& wings cut (T1)	766	540	1306	86.72	200	13.28	1506	261
Legs & wings cut (T2)	539	363	902	86.15	145	13.85	1047	180
Antenna, wings & legs cut (T3)	1015	139	1154	89.53	135	10.47	1289	231
Normal (T0)	682	225	907	87.97	124	12.03	1031	181

It is obvious from the outcomes that in treated clumps T1 and T3, where removal of antennae included, the eggs laid were additionally showing significance of antenna in egg lying when contrasted with different clusters where Antenna removal was not included (T2 and T0). Morohoshi (2000) detailed that among the four gatherings of secretary cells horizontally

situated in the rear mind, the IV bunch comprises of 3 Neuro-secretary cells are straightforwardly connected to the antennae on the head and their emission goes through external axon and arrives at the corpus cardiacum where it invigorates the organs and manages its capacity.

Table.4 Effect of treated eggs on hatching in Tasar silkworm

Treatments	Bivoltine eggs / g.95		Trivoltine eggs /g.85	
	Eggs kept	Hatching %	Eggs kept	Hatching %
Antenna & wings cut (T1)	475	91.25	425	91.00
Legs and wings cut (T2)	475	91.40	425	90.00
Antenna, wings & legs cut (T3)	475	88.95	425	89.00
Normal (T0)	475	90.20	425	90.00

6. Conclusion

It is inferred that when stress was given as body parts removal, the Neurohormones may offer sign to the cerebrum and invigorate to lay whatever eggs stayed in the mid-region. This sort of stress conditions referenced above might be worked on during grainage tasks of Tasar silk moth for improved egg laying and fruitfulness. The loads of moths, both genders, indicated abatement in outside worms aside from in the subsequent yield. Light of cases during diapause demonstrated a portion reliant, huge decrease with 5, 7.5 and

10 Gy, in the pupal span and increment in the length of the moth stage. Impact on silk fiber quality and structure are additionally noted with radiation. The positive improvement of a portion of the quantitative characters observed in the investigation shows the conceivable heterosis impact due to between ecorace crossings. Additionally, changeability in the tasar silkworm populaces could be of monstrous application to abuse or to exhibit heterosis impact in securing significant quantitative and subjective traits.

References

- Pandey, J.P, P.K.Mishra, D.Kumar, B.M. K. Singh and B.C.Prasad (2010) Effect of Temperature on Haemocytic immune responses of tropical Tasar Silkworm *Antheraea mylitta* D. Research.J.Immunol., 3: 169-177.
- Vemananda Reddy;G., Jayappa, T and Ravindra Singh (2016) Studies on egg production and egg handling techniques in Tasar Silkworm *Antheraea mylitta*. D. Annual report SSTL,(2015-16),SSTL, Central silk Board, Bangalore, India.
- Sudhakara Rao, P., K. M. Vijaya Kumari, B. Sailaja, G. V. Vishaka, S. Vidyunmala and Mishra, R. K. 2019. Studies on Egg Laying Behaviour and Fecundity under Different Stress Conditions in Tropical Tasar Silk Moth. *Int.J.Curr.Microbiol.App.Sci.* 8(09): 1182-1187.
- Barsagade D.D. and Tembhare B., Haemolymph amino acids and protein profile in the tropical Tasar silkworm, *Antheraea mylitta* (Drury) (Lepidoptera: Saturniidae) *Entomon*, 29(3), 261-266 (2004)
- Anita Singh, Ratnesh Kr Sharma, Bechan Sharma, Low temperature induced alterations in certain biochemical constituents of 5th instar larvae of *Philosamia ricini* (Lepidoptera: Saturniidae), *Open Access Insect Physiology*, 2, 11-16 (2010)
- Shamitha G. and Purushotham Rao, A Estimation of amino acids, urea, uric acid in tasar silkworm *Antheraea mylitta* D., *Journal of Environmental Biology*, 29(6), 893-896 (2008)
- Prashant, K.; Sharma, K.B. and Ali, Sarfaraz (2017) : Studies on the relative coupling behaviour of different tasar silkworms under various conditions. *Int. Journ of current research* (9): 54384-54387.
- Lokesh, G.; Srivastava, A.K.; Kar, P.K.; Srivastava, P.P. and Satay, A. (2016): Seasonal and climatic influence on the leaf biochemical of sal (*Shorea robusta*) flora and in situ breeding of Laria eco-race of *Antheraea mylitta* D. *Journ. of Entomology and Zoology studies.* 4 (6) 57-62.
- Srivastava (2011): Relationship of moth colour, fecundity and hatching in tropical tasar silkworm, *Antheraea mylitta* D. *Journ. Biospectra*, pp. 97-100.
- Ali, Sarfaraz & Sharma, K.B.. (2013). Impact of Environmental Factors on the Indoor Rearing Performances of Tropical Tasar Silkworm, *Antheraea mylitta* D.(Saturniidae: Lepidoptera). *The Ecoscan.* 4. 255-260.